

From Quantum Optics to Non-Commutative Geometry : A Non-Commutative Version of the Hopf Bundle, Veronese Mapping and Spin Representation

Kazuyuki FUJII *

Department of Mathematical Sciences

Yokohama City University

Yokohama, 236-0027

Japan

Abstract

In this paper we construct a non-commutative version of the Hopf bundle by making use of Jaynes-Comings model and so-called Quantum Diagonalization Method. The bundle has a kind of Dirac strings. However, they appear in only states containing the ground one $(\mathcal{F} \times \{|0\rangle\}) \cup \{|0\rangle\} \times \mathcal{F} \subset \mathcal{F} \times \mathcal{F}$ and don't appear in remaining excited states. This means that classical singularities are not universal in the process of non-commutativization.

Based on this construction we moreover give a non-commutative version of both the Veronese mapping which is the mapping from $\mathbf{CP}^1$ to $\mathbf{CP}^n$ with mapping degree n and the spin representation of the group $SU(2)$.

We also present some challenging problems concerning how classical (beautiful) properties can be extended to the non-commutative case.

*E-mail address : fujii@yokohama-cu.ac.jp

1 Introduction

This paper is an extended version of [1].

The Hopf bundles (which are famous examples of fiber bundles) over $\mathbf{K} = \mathbf{R}, \mathbf{C}, \mathbf{H}$ (the field of quaternion numbers), $\mathbf{O}$ (the field of octanion numbers) are classical objects and they are never written down in a local manner. If we write them locally then we are forced to encounter singular lines called the Dirac strings, see [1], [2].

It is very interesting to comment that the Hopf bundles correspond to topological solitons called Kink, Monopole, Instanton, Generalized Instanton respectively, see for example [2], [3], [4]. Therefore they are very important objects to study in detail.

Berry has given another expression to the Hopf bundle and Dirac strings by making use of a Hamiltonian (a simple spin model including the parameters x, y and z), see the paper(s) in [5]. We call this the Berry model for simplicity. In the following let us restrict to the case of $\mathbf{K}=\mathbf{C}$.

We would like to make the Hopf bundle non-commutative. Whether such a generalization is meaningful or not is not clear at the current time, however it may be worth trying, see for example [6] or more recently [7] and its references.

By the way, we are studying a quantum computation based on Cavity QED and one of the basic tools is the Jaynes–Cummings model (or more generally the Tavis–Cummings one), [8], [9], [10], [11]. This is given as a “half” of the Dicke model under the resonance condition and rotating wave approximation associated to it. If the resonance condition is not taken, then this model gives a non-commutative version of the Berry model. However, this new one is different from usual one because x and y coordinates are quantized, while z coordinate is not.

From the non-commutative Berry model we construct a non-commutative version of the Hopf bundle by making use of so-called Quantum Diagonalization Method (QDM) developed in [12]. Then we see that the Dirac strings appear in only states containing the ground one $(\mathcal{F} \times \{|0\rangle\} \cup \{|0\rangle\} \times \mathcal{F})$ where $\mathcal{F}$ is the Fock space generated by $\{a, a^\dagger, N = a^\dagger a\}$, while they don't appear in excited states $(\mathcal{F} \times \mathcal{F} - \mathcal{F} \times \{|0\rangle\} \cup \{|0\rangle\} \times \mathcal{F})$.

This means that classical singularities are not universal in the process of non-commutativization,

which is a very interesting phenomenon. This is one of reasons why we consider non-commutative generalizations (which are not necessarily unique) of classical geometry.

Moreover, we construct a non-commutative version of the Veronese mapping which is the mapping from $\mathbf{CP}^1$ to $\mathbf{CP}^n$ with mapping degree n . The mapping degree is usually defined by making use of the (first-) Chern class, so our mapping will become important if a non-commutative (or quantum) “Chern class” would be constructed.

We also challenge to construct a non-commutative version of the spin representation of group $SU(2)$. However, our trial is not enough because we could construct only the cases for spin $j = 1$ and $j = 3/2$. In this problem, we meet a very difficulty arising from the non-commutativity. Further study constructing a general theory will be required.

Why do we consider non-commutative versions of classical field models ? What is an advantage to consider such a generalization ? Researchers in this subject should answer such natural questions. This paper may give one of answers.

The contents of the paper are as follows :

Section 1 Introduction

Section 2 Berry Model and Dirac Strings : Review

Section 3 Non-Commutative Berry Model Arising from the Jaynes-Cummings Model

Section 4 Non-Commutative Hopf Bundle

Section 5 Non-Commutative Veronese Mapping

Section 6 Non-Commutative Representation Theory

Section 7 Discussion

Appendix

A Classical Theory of Projective Spaces

B Local Coordinate of the Projector

C Difficulty of Tensor Decomposition

2 Berry Model and Dirac Strings : Review

First of all we explain the Dirac strings and Hopf bundle which Berry constructed in [5]. The Hamiltonian considered by Berry is a simple spin model

$$H_B = x\sigma_1 + y\sigma_2 + z\sigma_3 = (x - iy)\sigma_+ + (x + iy)\sigma_- + z\sigma_3 = \begin{pmatrix} z & x - iy \\ x + iy & -z \end{pmatrix} \quad (1)$$

where σ_j ($j = 1 \sim 3$) is the Pauli matrices, $\sigma_{\pm} \equiv (1/2)(\sigma_1 \pm i\sigma_2)$ and x, y and z are parameters. We would like to diagonalize H_B above. The eigenvalues are

$$\lambda = \pm r \equiv \pm \sqrt{x^2 + y^2 + z^2}$$

and corresponding orthonormal eigenvectors are

$$|r\rangle = \frac{1}{\sqrt{2r(r+z)}} \begin{pmatrix} r+z \\ x+iy \end{pmatrix}, \quad |-r\rangle = \frac{1}{\sqrt{2r(r+z)}} \begin{pmatrix} -x+iy \\ r+z \end{pmatrix}.$$

Here we assume $(x, y, z) \in \mathbf{R}^3 - \{(0, 0, 0)\} \equiv \mathbf{R}^3 \setminus \{0\}$ to avoid a degenerate case. Therefore a unitary matrix defined by

$$U_I = (|r\rangle, |-r\rangle) = \frac{1}{\sqrt{2r(r+z)}} \begin{pmatrix} r+z & -x+iy \\ x+iy & r+z \end{pmatrix} \quad (2)$$

makes H_B diagonal like

$$H_B = U_I \begin{pmatrix} r & \\ & -r \end{pmatrix} U_I^\dagger \equiv U_I D_B U_I^\dagger. \quad (3)$$

We note that the unitary matrix U_I is not defined on the whole space $\mathbf{R}^3 \setminus \{0\}$. The defining region of U_I is

$$D_I = \mathbf{R}^3 \setminus \{0\} - \{(0, 0, z) \in \mathbf{R}^3 \mid z < 0\}. \quad (4)$$

The removed line $\{(0, 0, z) \in \mathbf{R}^3 \mid z < 0\}$ is just the (lower) Dirac string, which is impossible to add to D_I .

Next, we have another diagonal form of H_B like

$$H_B = U_{II} D_B U_{II}^\dagger \quad (5)$$




Figure 1: Dirac strings corresponding to I and II

with the unitary matrix U_{II} defined by

$$U_{II} = \frac{1}{\sqrt{2r(r-z)}} \begin{pmatrix} x - iy & -r + z \\ r - z & x + iy \end{pmatrix}. \quad (6)$$

The defining region of U_{II} is

$$D_{II} = \mathbf{R}^3 \setminus \{0\} - \{(0, 0, z) \in \mathbf{R}^3 \mid z > 0\}. \quad (7)$$

The removed line $\{(0, 0, z) \in \mathbf{R}^3 \mid z > 0\}$ is the (upper) Dirac string, which is also impossible to add to D_{II} .

Here we have diagonalizations of two types for H_B , so a natural question comes about. What is a relation between U_I and U_{II} ? If we define

$$\Phi = \frac{1}{\sqrt{x^2 + y^2}} \begin{pmatrix} x - iy & \\ & x + iy \end{pmatrix} \quad (8)$$

then it is easy to see

$$U_{II} = U_I \Phi.$$

We note that Φ (which is called a transition function) is not defined on the whole z -axis.

What we would like to emphasize here is that the diagonalization of H_B is not given globally (on $\mathbf{R}^3 \setminus \{0\}$). However, the dynamics is perfectly controlled by the system

$$\{(U_I, D_I), (U_{II}, D_{II}), \Phi, D_I \cup D_{II} = \mathbf{R}^3 \setminus \{0\}\}, \quad (9)$$

which defines a famous fiber bundle called the Hopf bundle associated to the complex numbers $\mathbf{C}$ ¹,

$$S^1 \longrightarrow S^3 \longrightarrow S^2,$$

see [2].

The projector corresponding to the Hopf bundle is given as

$$P(x, y, z) = U_I P_0 U_I^\dagger = U_{II} P_0 U_{II}^\dagger = \frac{1}{2r} \begin{pmatrix} r+z & x-iy \\ x+iy & r-z \end{pmatrix}, \quad (10)$$

where P_0 is a basic one

$$P_0 = \begin{pmatrix} 1 & \\ & 0 \end{pmatrix} \in M(2, \mathbf{C}).$$

We note that in (10) Dirac strings don't appear because the projector P is expressed globally.

3 Non-Commutative Berry Model Arising from the Jaynes-Cummings Model

First, let us explain the Jaynes-Cummings model which is well-known in quantum optics, [8], [9]. The Hamiltonian of Jaynes-Cummings model can be written as follows (we set $\hbar = 1$ for simplicity)

$$H = \omega 1_2 \otimes a^\dagger a + \frac{\Delta}{2} \sigma_3 \otimes \mathbf{1} + g (\sigma_+ \otimes a + \sigma_- \otimes a^\dagger), \quad (11)$$

where ω is the frequency of single radiation field, Δ the energy difference of two level atom, a and $a^\dagger$ are annihilation and creation operators of the field, and g a coupling constant. We assume that g is small enough (a weak coupling regime). Here σ_+ , σ_- and σ_3 are given as

$$\sigma_+ = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \sigma_- = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad 1_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}. \quad (12)$$

See the figure 2 as an image of the Jaynes-Cummings model (we don't repeat here).

¹The base space $\mathbf{R}^3 \setminus \{0\}$ is homotopic to the two-dimensional sphere S^2




Figure 2: One atom and a single photon inserted in a cavity

Now we consider the evolution operator of the model. We rewrite the Hamiltonian (11) as follows.

$$H = \omega 1_2 \otimes a^\dagger a + \frac{\omega}{2} \sigma_3 \otimes \mathbf{1} + \frac{\Delta - \omega}{2} \sigma_3 \otimes \mathbf{1} + g (\sigma_+ \otimes a + \sigma_- \otimes a^\dagger) \equiv H_1 + H_2. \quad (13)$$

Then it is easy to see $[H_1, H_2] = 0$, which leads to $e^{-itH} = e^{-itH_1} e^{-itH_2}$.

In the following we consider e^{-itH_2} in which the resonance condition $\Delta - \omega = 0$ is not taken.

For simplicity we set $\theta = \frac{\Delta - \omega}{2g} (\neq 0)$ ² then

$$H_2 = g \left(\sigma_+ \otimes a + \sigma_- \otimes a^\dagger + \frac{\Delta - \omega}{2g} \sigma_3 \otimes \mathbf{1} \right) = g \left(\sigma_+ \otimes a + \sigma_- \otimes a^\dagger + \theta \sigma_3 \otimes \mathbf{1} \right).$$

For further simplicity we set

$$H_{JC} = \sigma_+ \otimes a + \sigma_- \otimes a^\dagger + \theta \sigma_3 \otimes \mathbf{1} = \begin{pmatrix} \theta & a \\ a^\dagger & -\theta \end{pmatrix}, \quad [a, a^\dagger] = \mathbf{1} \quad (14)$$

where we have written θ in place of $\theta \mathbf{1}$ for simplicity.

H_{JC} can be considered as a non-commutative version of H_B under the quantum-classical correspondence $a \longleftrightarrow x - iy$, $a^\dagger \longleftrightarrow x + iy$ and $\theta \longleftrightarrow z$:

$$H_B = \begin{pmatrix} z & x - iy \\ x + iy & -z \end{pmatrix}, \quad [x - iy, x + iy] = 0 \quad \longrightarrow \quad H_{JC} = \begin{pmatrix} \theta & a \\ a^\dagger & -\theta \end{pmatrix}, \quad [a, a^\dagger] = \mathbf{1}.$$

That is, x and y coordinates are quantized, while z coordinate is not, which is different from usual one, see for example [6]. It may be possible for us to call this **a non-commutative Berry model**. We note that this model is derived not “by hand” but by the model in quantum optics itself.

²Since the Jaynes–Cummings model is obtained by the Dicke model under some resonance condition on parameters included, it is nothing but an approximate one in the neighborhood of the point, so we must assume that $|\theta|$ is small enough. However, as a model in mathematical physics there is no problem to take θ be arbitrary

4 Non-Commutative Hopf Bundle

We usually analyze (14) by reducing it to each component contained in $H(2, \mathbf{C})$, which is a typical analytic method. However, we don't adopt such a method.

First we make the Hamiltonian (14) diagonal like in Section 2 and research whether “Dirac strings” exist or not in this non-commutative model, which is very interesting from not only quantum optical but also mathematical point of view.

It is easy to see

$$H_{JC} = \begin{pmatrix} \theta & a \\ a^\dagger & -\theta \end{pmatrix} = \begin{pmatrix} 1 & \\ & a^\dagger \frac{1}{\sqrt{N+1}} \end{pmatrix} \begin{pmatrix} \theta & \sqrt{N+1} \\ \sqrt{N+1} & -\theta \end{pmatrix} \begin{pmatrix} 1 & \\ & \frac{1}{\sqrt{N+1}}a \end{pmatrix} \quad (15)$$

from [12], where N is the number operator $N = a^\dagger a$. Then the middle matrix in the right hand side can be considered as a classical one, so we can diagonalize it easily

$$\begin{pmatrix} \theta & \sqrt{N+1} \\ \sqrt{N+1} & -\theta \end{pmatrix} = \begin{cases} U_I \begin{pmatrix} R(N+1) & \\ & -R(N+1) \end{pmatrix} U_I^\dagger \\ U_{II} \begin{pmatrix} R(N+1) & \\ & -R(N+1) \end{pmatrix} U_{II}^\dagger \end{cases} \quad (16)$$

where

$$R(N) = \sqrt{N + \theta^2}$$

and U_I, U_{II} are given by

$$U_I = \frac{1}{\sqrt{2R(N+1)(R(N+1) + \theta)}} \begin{pmatrix} R(N+1) + \theta & -\sqrt{N+1} \\ \sqrt{N+1} & R(N+1) + \theta \end{pmatrix}, \quad (17)$$

$$U_{II} = \frac{1}{\sqrt{2R(N+1)(R(N+1) - \theta)}} \begin{pmatrix} \sqrt{N+1} & -R(N+1) + \theta \\ R(N+1) - \theta & \sqrt{N+1} \end{pmatrix}. \quad (18)$$

Now let us rewrite (15) by making use of (16) with (17). Inserting the identity

$$\begin{pmatrix} 1 & \\ & \frac{1}{\sqrt{N+1}}a \end{pmatrix} \begin{pmatrix} 1 & \\ & a^\dagger \frac{1}{\sqrt{N+1}} \end{pmatrix} = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$$

gives

$$\begin{aligned}
H_{JC} &= \begin{pmatrix} 1 & \\ & a^\dagger \frac{1}{\sqrt{N+1}} \end{pmatrix} U_I \begin{pmatrix} R(N+1) & \\ & -R(N+1) \end{pmatrix} U_I^\dagger \begin{pmatrix} 1 & \\ & \frac{1}{\sqrt{N+1}} a \end{pmatrix} \\
&= \begin{pmatrix} 1 & \\ & a^\dagger \frac{1}{\sqrt{N+1}} \end{pmatrix} U_I \begin{pmatrix} 1 & \\ & \frac{1}{\sqrt{N+1}} a \end{pmatrix} \begin{pmatrix} 1 & \\ & a^\dagger \frac{1}{\sqrt{N+1}} \end{pmatrix} \begin{pmatrix} R(N+1) & \\ & -R(N+1) \end{pmatrix} \times \\
&\quad \begin{pmatrix} 1 & \\ & \frac{1}{\sqrt{N+1}} a \end{pmatrix} \begin{pmatrix} 1 & \\ & a^\dagger \frac{1}{\sqrt{N+1}} \end{pmatrix} U_I^\dagger \begin{pmatrix} 1 & \\ & \frac{1}{\sqrt{N+1}} a \end{pmatrix} \\
&= V_I \begin{pmatrix} R(N+1) & \\ & -R(N) \end{pmatrix} V_I^\dagger, \tag{19}
\end{aligned}$$

where

$$\begin{aligned}
V_I &= \begin{pmatrix} \frac{1}{\sqrt{2R(N+1)(R(N+1)+\theta)}} & \\ & \frac{1}{\sqrt{2R(N)(R(N)+\theta)}} \end{pmatrix} \begin{pmatrix} R(N+1)+\theta & -a \\ & a^\dagger \end{pmatrix} \\
&= \begin{pmatrix} R(N+1)+\theta & -a \\ & a^\dagger \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{2R(N+1)(R(N+1)+\theta)}} & \\ & \frac{1}{\sqrt{2R(N)(R(N)+\theta)}} \end{pmatrix}. \tag{20}
\end{aligned}$$

Similarly, we can rewrite (15) by making use of (16) with (18). By inserting the identity

$$\begin{pmatrix} \frac{1}{\sqrt{N+1}} a & \\ & 1 \end{pmatrix} \begin{pmatrix} a^\dagger \frac{1}{\sqrt{N+1}} & \\ & 1 \end{pmatrix} = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$$

we obtain

$$H_{JC} = V_{II} \begin{pmatrix} R(N) & \\ & -R(N+1) \end{pmatrix} V_{II}^\dagger, \tag{21}$$

where

$$\begin{aligned}
V_{II} &= \begin{pmatrix} \frac{1}{\sqrt{2R(N+1)(R(N+1)-\theta)}} & \\ & \frac{1}{\sqrt{2R(N)(R(N)-\theta)}} \end{pmatrix} \begin{pmatrix} a & -R(N+1)+\theta \\ R(N)-\theta & a^\dagger \end{pmatrix} \\
&= \begin{pmatrix} a & -R(N+1)+\theta \\ R(N)-\theta & a^\dagger \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{2R(N)(R(N)-\theta)}} & \\ & \frac{1}{\sqrt{2R(N+1)(R(N+1)-\theta)}} \end{pmatrix}. \tag{22}
\end{aligned}$$

Tidying up these we have

$$H_{JC} = \begin{cases} V_I \begin{pmatrix} R(N+1) & \\ & -R(N) \end{pmatrix} V_I^\dagger \\ V_{II} \begin{pmatrix} R(N) & \\ & -R(N+1) \end{pmatrix} V_{II}^\dagger \end{cases} \quad (23)$$

with V_I and V_{II} above. From the equations

$$R(N+1)|0\rangle = \sqrt{1+\theta^2} > \theta, \quad R(N)|0\rangle = \sqrt{\theta^2} = |\theta|$$

we know

$$(R(N) \pm \theta) |0\rangle = (|\theta| \pm \theta) |0\rangle,$$

so the strings corresponding to Dirac ones exist in only states $\mathcal{F} \times \{|0\rangle\} \cup \{|0\rangle\} \times \mathcal{F}$ where $\mathcal{F}$ is the Fock space generated by $\{a, a^\dagger, N\}$, while in other excited states $\mathcal{F} \times \mathcal{F} \setminus \mathcal{F} \times \{|0\rangle\} \cup \{|0\rangle\} \times \mathcal{F}$ they don't exist³, see the figure 3. The phenomenon is very interesting. For simplicity we again call these strings Dirac ones in the following.

The “parameter space” of H_{JC} can be identified with $\mathcal{F} \times \mathcal{F} \times \mathbf{R} \ni (*, *, \theta)$, so the domains D_I of V_I and D_{II} of V_{II} are respectively

$$D_I = \mathcal{F} \times \mathcal{F} \times \mathbf{R} - \mathcal{F} \times \{|0\rangle\} \times \mathbf{R}_{\leq 0}, \quad (24)$$

$$D_{II} = \mathcal{F} \times \mathcal{F} \times \mathbf{R} - (\mathcal{F} \times \{|0\rangle\} \cup \{|0\rangle\} \times \mathcal{F}) \times \mathbf{R}_{\geq 0} \quad (25)$$

by (20) and (22). We note that

$$D_I \cup D_{II} = \mathcal{F} \times \mathcal{F} \times \mathbf{R} - \mathcal{F} \times \{|0\rangle\} \times \{\theta = 0\}.$$

Then the transition “function” (operator) is given by

$$\Phi_{JC} = \begin{pmatrix} a \frac{1}{\sqrt{N}} & \\ & \frac{1}{\sqrt{N}} a^\dagger \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{N+1}} a & \\ & a^\dagger \frac{1}{\sqrt{N+1}} \end{pmatrix}.$$

Therefore the system

$$\{(V_I, D_I), (V_{II}, D_{II}), \Phi_{JC}, D_I \cup D_{II}\} \quad (26)$$

³We have identified $\mathcal{F} \times \mathcal{F}$ with the space of 2-component vectors over $\mathcal{F}$




Figure 3: The bases of $\mathcal{F} \times \mathcal{F}$. The black circle means bases giving Dirac strings, while the white one don't.

is a non-commutative version of the Hopf bundle (9). The projector in this case becomes

$$\begin{aligned}
 P_{JC} &= V_I \begin{pmatrix} \mathbf{1} & \\ & \mathbf{0} \end{pmatrix} V_I^\dagger = V_{II} \begin{pmatrix} \mathbf{1} & \\ & \mathbf{0} \end{pmatrix} V_{II}^\dagger \\
 &= \begin{cases} \begin{pmatrix} \frac{1}{2R(N+1)} & \\ & \frac{1}{2R(N)} \end{pmatrix} \begin{pmatrix} R(N+1) + \theta & a \\ a^\dagger & R(N) - \theta \end{pmatrix} \\ \begin{pmatrix} R(N+1) + \theta & a \\ a^\dagger & R(N) - \theta \end{pmatrix} \begin{pmatrix} \frac{1}{2R(N+1)} & \\ & \frac{1}{2R(N)} \end{pmatrix} \end{cases}. \quad (27)
 \end{aligned}$$

Note that the projector P_{JC} is not defined on $\mathcal{F} \times \{|0\rangle\} \times \{\theta = 0\} = \mathcal{F} \times \mathcal{F} \times \mathbf{R} - D_I \cup D_{II}$.

A comment is in order. From (27) we obtain a quantum version of (classical) spectral decomposition (a “quantum spectral decomposition” by Suzuki [13])

$$H_{JC} = \begin{pmatrix} R(N+1) & \\ & R(N) \end{pmatrix} P_{JC} - \begin{pmatrix} R(N+1) & \\ & R(N) \end{pmatrix} (\mathbf{1}_2 - P_{JC}). \quad (28)$$

As a bonus of the decomposition let us rederive the calculation of $e^{-igtH_{JC}}$ which has been

given in [9]. The result is

$$e^{-igtH_{JC}} = \begin{pmatrix} \cos(tgR(N+1)) - i\theta \frac{\sin(tgR(N+1))}{R(N+1)} & -i \frac{\sin(tgR(N+1))}{R(N+1)} a \\ -i \frac{\sin(tgR(N))}{R(N)} a^\dagger & \cos(tgR(N)) + i\theta \frac{\sin(tgR(N))}{R(N)} \end{pmatrix} \quad (29)$$

by making use of (23) (or (28)). We leave it to the readers.

5 Non-Commutative Veronese Mapping

Let us make a brief review of the Veronese mapping. The map

$$\mathbf{CP}^1 \longrightarrow \mathbf{CP}^n$$

is defined as

$$[z_1 : z_2] \longrightarrow \left[z_1^n : \sqrt{n C_1} z_1^{n-1} z_2 : \cdots : \sqrt{n C_j} z_1^{n-j} z_2^j : \cdots : \sqrt{n C_{n-1}} z_1 z_2^{n-1} : z_2^n \right]$$

by making use of the homogeneous coordinate, see Appendix A. We also have another expression of this map by using

$$S_{\mathbf{C}}^1 \longrightarrow S_{\mathbf{C}}^n : v_1 \equiv \begin{pmatrix} z_1 \\ z_2 \end{pmatrix} \longrightarrow v_n \equiv \begin{pmatrix} z_1^n \\ \sqrt{n C_1} z_1^{n-1} z_2 \\ \vdots \\ \sqrt{n C_j} z_1^{n-j} z_2^j \\ \vdots \\ \sqrt{n C_{n-1}} z_1 z_2^{n-1} \\ z_2^n \end{pmatrix}, \quad |z_1|^2 + |z_2|^2 = 1$$

where $S_{\mathbf{C}}^m = \left\{ (w_1, w_2, \dots, w_{m+1})^T \in \mathbf{C}^{m+1} \mid \sum_{j=1}^{m+1} |w_j|^2 = 1 \right\} \cong S^{2m+1}$ and $\mathbf{CP}^m = S_{\mathbf{C}}^m / U(1)$.

Then the Veronese mapping is also written as

$$\mathbf{CP}^1 \longrightarrow \mathbf{CP}^n : P_1 = v_1 v_1^\dagger \longmapsto P_n = v_n v_n^\dagger.$$

by using projectors.

Moreover, the local map ($z \equiv z_2/z_1$) is given as

$$\mathbf{C} \longrightarrow \mathbf{C}^n : z \longrightarrow \begin{pmatrix} \sqrt{n C_1} z \\ \vdots \\ \sqrt{n C_j} z^j \\ \vdots \\ \sqrt{n C_{n-1}} z^{n-1} \\ z^n \end{pmatrix}.$$

See the following picture as a whole.

$$\begin{array}{ccc} S_{\mathbf{C}}^1 & \longrightarrow & S_{\mathbf{C}}^n \\ \downarrow & & \downarrow \\ \mathbf{C}P^1 & \longrightarrow & \mathbf{C}P^n \\ \uparrow & & \uparrow \\ \mathbf{C} & \longrightarrow & \mathbf{C}^n \end{array}$$

Next we want to consider a non-commutative version of the map. In the following we treat vectors as column ones. If we set

$$\mathcal{A} \equiv \begin{pmatrix} X_0 \\ Y_0 \end{pmatrix} = \begin{pmatrix} \frac{R(N+1)+\theta}{\sqrt{2R(N+1)(R(N+1)+\theta)}} \\ \frac{1}{\sqrt{2R(N)(R(N)+\theta)}} a^\dagger \end{pmatrix} \quad (30)$$

from V_I in (20), then

$$\mathcal{A}^\dagger \mathcal{A} = X_0^2 + Y_0^\dagger Y_0 = \mathbf{1} \quad \text{and} \quad Y_0 X_0^{-1} = \frac{1}{R(N) + \theta} a^\dagger = Z.$$

That is, $\mathcal{A} = (X_0, Y_0)^T$ is a non-commutative “sphere” and Z is a kind of “stereographic projection” of the sphere. It is easy to see

$$\mathbf{1} + Z^\dagger Z = \frac{2R(N+1)}{R(N+1) + \theta} = X_0^{-2} \implies X_0 = (\mathbf{1} + Z^\dagger Z)^{-1/2}. \quad (31)$$

Here let us introduce new notations for the following. For $j \geq 0$ we set

$$X_{-j} = \frac{R(N+1-j) + \theta}{\sqrt{2R(N+1-j)(R(N+1-j) + \theta)}}, \quad (32)$$

$$Y_{-j} = \sqrt{\frac{N-j}{N}} \frac{1}{\sqrt{2R(N-j)(R(N-j) + \theta)}} a^\dagger. \quad (33)$$

We list some useful formulas.

$$X_{-j}^2 + Y_{-j}^\dagger Y_{-j} = \mathbf{1} \quad \text{and} \quad Y_{-j}^\dagger Y_{-j} = Y_{-j+1} Y_{-j+1}^\dagger \quad \text{for } j \geq 0. \quad (34)$$

Now we are in a position to define a quantum version of the Veronese mapping which plays a very important role in “classical” Mathematics.

$$\mathcal{A} = \begin{pmatrix} X_0 \\ Y_0 \end{pmatrix} \longrightarrow \mathcal{A}_n = \begin{pmatrix} X_0^n \\ \sqrt{n} C_1 Y_0 X_0^{n-1} \\ \vdots \\ \sqrt{n} C_j Y_{-(j-1)} Y_{-(j-2)} \cdots Y_{-1} Y_0 X_0^{n-j} \\ \vdots \\ \sqrt{n} C_{n-1} Y_{-(n-2)} Y_{-(n-3)} \cdots Y_{-1} Y_0 X_0 \\ Y_{-(n-1)} Y_{-(n-2)} Y_{-(n-3)} \cdots Y_{-1} Y_0 \end{pmatrix}. \quad (35)$$

Then it is not difficult to see

$$\mathcal{A}_n^\dagger \mathcal{A}_n = (X_0^2 + Y_0^\dagger Y_0)^n = (\mathcal{A}^\dagger \mathcal{A})^n = \mathbf{1}.$$

From this we can define the projectors which correspond to projective spaces like

$$\mathcal{P}_n = \mathcal{A}_n \mathcal{A}_n^\dagger, \quad \mathcal{P}_1 = \mathcal{A} \mathcal{A}^\dagger, \quad (36)$$

so the map

$$\mathcal{P}_1 \longrightarrow \mathcal{P}_n \quad (37)$$

is a non-commutative version of the Veronese mapping.

Next, we define a local “coordinate” of the Veronese mapping defined above.

$$\begin{aligned}
\mathcal{A}_n &= \begin{pmatrix} \mathbf{1} \\ \sqrt{nC_1}Y_0X_0^{-1} \\ \vdots \\ \sqrt{nC_j}Y_{-(j-1)}Y_{-(j-2)}\cdots Y_{-1}Y_0X_0^{-j} \\ \vdots \\ \sqrt{nC_{n-1}}Y_{-(n-2)}Y_{-(n-3)}\cdots Y_{-1}Y_0X_0^{-(n-1)} \\ Y_{-(n-1)}Y_{-(n-2)}Y_{-(n-3)}\cdots Y_{-1}Y_0X_0^{-n} \end{pmatrix} X_0^n \\
&= \dots \\
&= \begin{pmatrix} \mathbf{1} \\ \sqrt{nC_1}Y_0X_0^{-1} \\ \vdots \\ \sqrt{nC_j}Y_{-(j-1)}X_{-(j-1)}^{-1}Y_{-(j-2)}X_{-(j-2)}^{-1}\cdots Y_{-1}X_{-1}^{-1}Y_0X_0^{-1} \\ \vdots \\ \sqrt{nC_{n-1}}Y_{-(n-2)}X_{-(n-2)}^{-1}Y_{-(n-3)}X_{-(n-3)}^{-1}\cdots Y_{-1}X_{-1}^{-1}Y_0X_0^{-1} \\ Y_{-(n-1)}X_{-(n-1)}^{-1}Y_{-(n-2)}X_{-(n-2)}^{-1}Y_{-(n-3)}X_{-(n-3)}^{-1}\cdots Y_{-1}X_{-1}^{-1}Y_0X_0^{-1} \end{pmatrix} X_0^n
\end{aligned}$$

where we have used the relation

$$Y_{-j}X_{-k}^{-1} = X_{-(k+1)}^{-1}Y_{-j}$$

due to $a^\dagger$ in Y_{-j} . Moreover, by (32) and (33)

$$Y_{-j}X_{-j}^{-1} = \sqrt{\frac{N-j}{N}} \frac{1}{R(N-j) + \theta} a^\dagger \equiv Z_{-j} \quad \text{for } j \geq 0.$$

Note that $Z_0 = Z$. Therefore by using (31) we have

$$\mathcal{A}_n = \begin{pmatrix} \mathbf{1} \\ \sqrt{nC_1}Z_0 \\ \vdots \\ \sqrt{nC_j}Z_{-(j-1)}Z_{-(j-2)}\cdots Z_{-1}Z_0 \\ \vdots \\ \sqrt{nC_{n-1}}Z_{-(n-2)}Z_{-(n-3)}\cdots Z_{-1}Z_0 \\ Z_{-(n-1)}Z_{-(n-2)}Z_{-(n-3)}\cdots Z_{-1}Z_0 \end{pmatrix} (\mathbf{1} + Z_0^\dagger Z_0)^{-n/2}. \quad (38)$$

Now if we define

$$\mathcal{Z}_n = \begin{pmatrix} \sqrt{nC_1}Z_0 \\ \vdots \\ \sqrt{nC_j}Z_{-(j-1)}Z_{-(j-2)}\cdots Z_{-1}Z_0 \\ \vdots \\ \sqrt{nC_{n-1}}Z_{-(n-2)}Z_{-(n-3)}\cdots Z_{-1}Z_0 \\ Z_{-(n-1)}Z_{-(n-2)}Z_{-(n-3)}\cdots Z_{-1}Z_0 \end{pmatrix}, \quad (39)$$

then

$$\mathcal{A}_n = \begin{pmatrix} \mathbf{1} \\ \mathcal{Z}_n \end{pmatrix} (\mathbf{1} + Z_0^\dagger Z_0)^{-n/2}$$

and it is easy to show

$$\mathbf{1} + \mathcal{Z}_n^\dagger \mathcal{Z}_n = (\mathbf{1} + Z_0^\dagger Z_0)^n,$$

so we obtain

$$\begin{aligned} \mathcal{P}_n &= \mathcal{A}_n \mathcal{A}_n^\dagger \\ &= \begin{pmatrix} \mathbf{1} \\ \mathcal{Z}_n \end{pmatrix} (\mathbf{1} + \mathcal{Z}_n^\dagger \mathcal{Z}_n)^{-1} (\mathbf{1}, \mathcal{Z}_n^\dagger) \\ &= \begin{pmatrix} (\mathbf{1} + \mathcal{Z}_n^\dagger \mathcal{Z}_n)^{-1} & (\mathbf{1} + \mathcal{Z}_n^\dagger \mathcal{Z}_n)^{-1} \mathcal{Z}_n^\dagger \\ \mathcal{Z}_n (\mathbf{1} + \mathcal{Z}_n^\dagger \mathcal{Z}_n)^{-1} & \mathcal{Z}_n (\mathbf{1} + \mathcal{Z}_n^\dagger \mathcal{Z}_n)^{-1} \mathcal{Z}_n^\dagger \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{1} & -\mathcal{Z}_n^\dagger \\ \mathcal{Z}_n & \mathbf{1} \end{pmatrix} \begin{pmatrix} \mathbf{1} & \\ & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{1} & -\mathcal{Z}_n^\dagger \\ \mathcal{Z}_n & \mathbf{1} \end{pmatrix}^{-1}. \end{aligned} \quad (40)$$

This is the Oike expression in [14], see also Appendix B.

A comment is in order. Two of important properties which the classical Veronese mapping has are

1. The Veronese mapping $\mathbf{CP}^1 \longrightarrow \mathbf{CP}^n$ has the **mapping degree** n
2. The Veronese surface (which is the image of Veronese mapping) is a **minimal surface** in $\mathbf{CP}^n$

Since we have constructed a non-commutative version of the Veronese mapping, a natural question arises : What are non-commutative versions corresponding to 1. and 2. above ?

These are very interesting problems from the view point of non-commutative “differential” geometry. It is worth challenging.

6 Non-Commutative Representation Theory

The construction of spin j -representation ($j \in \mathbf{Z}_{\geq 0} + 1/2$) is very well-known. Let us make a brief review within our necessity. For the vector space

$$\mathcal{H}_J = \text{Vect}_{\mathbf{C}} \left\{ \frac{z^{j-m}}{\sqrt{(j-m)!(j+m)!}} \mid m \in \{-j, -j+1, \dots, j-1, j\} \right\}$$

where $J = 2j + 1 (\in \mathbf{N})$, the inner product in this space is given by

$$\left(\sum_{l=0}^{J-1} a_l z^l, \sum_{l=0}^{J-1} b_l z^l \right) = \sum_{l=0}^{J-1} l!(J-1-l)! a_l \bar{b}_l.$$

For example, for $j = 1/2$, $j = 1$ and $j = 3/2$

$$\mathcal{H}_2 = \text{Vect}_{\mathbf{C}} \{z, 1\}, \quad \mathcal{H}_3 = \text{Vect}_{\mathbf{C}} \left\{ \frac{z^2}{\sqrt{2}}, z, \frac{1}{\sqrt{2}} \right\}, \quad \mathcal{H}_4 = \text{Vect}_{\mathbf{C}} \left\{ \frac{z^3}{\sqrt{6}}, \frac{z^2}{\sqrt{2}}, \frac{z}{\sqrt{2}}, \frac{1}{\sqrt{6}} \right\}.$$

For

$$A = \begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix} \in SU(2) \quad (|\alpha|^2 + |\beta|^2 = 1)$$

the spin j representation

$$\phi_j : SU(2) \longrightarrow SU(J)$$

is defined as

$$(\phi_j(A)f)(z) = (-\bar{\beta}z + \bar{\alpha})^{j-1} f\left(\frac{\alpha z + \beta}{-\bar{\beta}z + \bar{\alpha}}\right) \quad (41)$$

where $f \in \mathcal{H}_J$. It is easy to obtain $\phi_j(A)$ for $j = 1/2$, $j = 1$ and $j = 3/2$.

Namely, the spin 1/2 representation is

$$\phi_{1/2}(A) = \begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix} = A, \quad (42)$$

the spin 1 representation is

$$\phi_1(A) = \begin{pmatrix} \alpha^2 & -\sqrt{2}\alpha\bar{\beta} & \bar{\beta}^2 \\ \sqrt{2}\alpha\beta & |\alpha|^2 - |\beta|^2 & -\sqrt{2}\bar{\alpha}\bar{\beta} \\ \beta^2 & \sqrt{2}\bar{\alpha}\beta & \bar{\alpha}^2 \end{pmatrix}, \quad (43)$$

and the spin 3/2 representation is

$$\phi_{3/2}(A) = \begin{pmatrix} \alpha^3 & -\sqrt{3}\alpha^2\bar{\beta} & \sqrt{3}\alpha\bar{\beta}^2 & -\bar{\beta}^3 \\ \sqrt{3}\alpha^2\beta & (|\alpha|^2 - 2|\beta|^2)\alpha & -(2|\alpha|^2 - |\beta|^2)\bar{\beta} & \sqrt{3}\bar{\alpha}\bar{\beta}^2 \\ \sqrt{3}\alpha\beta^2 & (2|\alpha|^2 - |\beta|^2)\beta & (|\alpha|^2 - 2|\beta|^2)\bar{\alpha} & -\sqrt{3}\bar{\alpha}^2\bar{\beta} \\ \beta^3 & \sqrt{3}\bar{\alpha}\beta^2 & \sqrt{3}\bar{\alpha}^2\beta & \bar{\alpha}^3 \end{pmatrix}. \quad (44)$$

Next we want to consider a non-commutative version of the spin representation. However, since such a theory has not been known as far as we know we must look for mappings corresponding to $\phi_1(A)$ and $\phi_{3/2}(A)$ by (many) trial and error, see Appendix C.

If we set

$$V \equiv V_I = \begin{pmatrix} X_0 & -Y_0^\dagger \\ Y_0 & X_{-1} \end{pmatrix} : \text{unitary}$$

from (20), then the corresponding map for $\phi_1(A)$ is

$$\Phi_1(V) = \begin{pmatrix} X_0^2 & -\sqrt{2}X_0Y_0^\dagger & Y_0^\dagger Y_{-1}^\dagger \\ \sqrt{2}Y_0X_0 & X_{-1}^2 - Y_{-1}^\dagger Y_{-1} & -\sqrt{2}X_{-1}Y_{-1}^\dagger \\ Y_{-1}Y_0 & \sqrt{2}Y_{-1}X_{-1} & X_{-2}^2 \end{pmatrix} \quad (45)$$

and the corresponding map for $\phi_{3/2}(A)$ is

$$\Phi_{3/2}(V) = \begin{pmatrix} X_0^3 & -\sqrt{3}X_0^2Y_0^\dagger & \sqrt{3}X_0Y_0^\dagger Y_{-1}^\dagger & -Y_0^\dagger Y_{-1}^\dagger Y_{-2}^\dagger \\ \sqrt{3}Y_0X_0^2 & X_{-1}(X_{-1}^2 - 2Y_{-1}^\dagger Y_{-1}) & -(2X_{-1}^2 - Y_{-1}^\dagger Y_{-1})Y_{-1}^\dagger & \sqrt{3}X_{-1}Y_{-1}^\dagger Y_{-2}^\dagger \\ \sqrt{3}Y_{-1}Y_0X_0 & Y_{-1}(2X_{-1}^2 - Y_{-1}^\dagger Y_{-1}) & X_{-2}(X_{-2}^2 - 2Y_{-2}^\dagger Y_{-2}) & -\sqrt{3}X_{-2}^2Y_{-2}^\dagger \\ Y_{-2}Y_{-1}Y_0 & \sqrt{3}Y_{-2}Y_{-1}X_{-1} & \sqrt{3}Y_{-2}X_{-2}^2 & X_{-3}^3 \end{pmatrix}. \quad (46)$$

To check the unitarity of $\Phi_1(V)$ and $\Phi_{3/2}(V)$ is long but straightforward.

For $j \geq 2$ we could not find a general method like (41) which determines $\Phi_j(V)$. However, we know only that the first column of $\Phi_j(V)$ is just $\mathcal{A}_{2j}$ in (35).,

$$\Phi_j(V) = (\mathcal{A}_{2j}, *, \dots, *) : \text{unitary}$$

and

$$\Phi_j(V) \begin{pmatrix} \mathbf{1} & & & \\ & \mathbf{0} & & \\ & & \ddots & \\ & & & \mathbf{0} \end{pmatrix} \Phi_j(V)^\dagger = \mathcal{A}_{2j}\mathcal{A}_{2j}^\dagger = \mathcal{P}_{2j}.$$

We leave finding a general method to the readers as a challenging problem.

7 Discussion

In this paper we derived a non-commutative version of the Berry model from the Jaynes–Cummings model in quantum optics and constructed a non-commutative version of the Hopf bundle.

The bundle has a kind of Dirac strings. However, they appear in only states containing the ground one ($\mathcal{F} \times \{|0\rangle\} \cup \{|0\rangle\} \times \mathcal{F} \subset \mathcal{F} \times \mathcal{F}$) and don't appear in excited states, which is very interesting.

In general, a non-commutative version of classical field theory is of course not unique. If our model is a “correct” one, then this paper give an example that classical singularities like Dirac

strings are not universal in some non-commutative model. As to general case with higher spins which are not easy see [13].

More generally, it is probable that a singularity (singularities) in some classical model is (are) removed in the process of non-commutativization.

Moreover, based on this model a non-commutative version of the Veronese mapping or spin representation was constructed. The results in the paper will become a starting point to construct fruitful non-commutative geometry.

Last, we would like to make a comment. To develop a “quantum” mathematics we need a rigorous method to treat an analysis or a geometry on infinite dimensional spaces like Fock space. In quantum field theories physicists have given some (interesting) methods, while they are more or less formal from the mathematical point of view. It is a rigorous method which we need. As a trial [16] is recommended.

Acknowledgment.

The author wishes to thank Akira Asada, Yoshinori Machida, Ryu Sasaki and Tatsuo Suzuki for their helpful comments and suggestions.

Appendix

A Classical Theory of Projective Spaces

Complex projective spaces are typical examples of symmetric spaces and are very tractable, so they are used to construct several examples in both physics and mathematics.

We make a review of complex projective spaces within our necessity, see for example [2], [14], [15].

For $n \in \mathbf{N}$ the complex projective space $\mathbf{CP}^n$ is defined as follows : For $\boldsymbol{\zeta}, \boldsymbol{\mu} \in \mathbf{C}^{n+1} - \{\mathbf{0}\}$ $\boldsymbol{\zeta}$ is equivalent to $\boldsymbol{\mu}$ ($\boldsymbol{\zeta} \sim \boldsymbol{\mu}$) if and only if $\boldsymbol{\zeta} = \lambda \boldsymbol{\mu}$ for $\lambda \in \mathbf{C} - \{0\}$. We show the equivalent relation class as $[\boldsymbol{\zeta}]$ and set $\mathbf{CP}^n \equiv \mathbf{C}^{n+1} - \{\mathbf{0}\} / \sim$. For $\boldsymbol{\zeta} = (\zeta_0, \zeta_1, \dots, \zeta_n)$ we write usually as

$[\zeta] = [\zeta_0 : \zeta_1 : \cdots : \zeta_n]$. Then it is well-known that $\mathbf{CP}^n$ has $n + 1$ local charts, namely

$$\mathbf{CP}^n = \bigcup_{j=0}^n U_j, \quad U_j = \{[\zeta_0 : \cdots : \zeta_j : \cdots : \zeta_n] \mid \zeta_j \neq 0\}. \quad (47)$$

Since

$$(\zeta_0, \dots, \zeta_j, \dots, \zeta_n) = \zeta_j \left(\frac{\zeta_0}{\zeta_j}, \dots, \frac{\zeta_{j-1}}{\zeta_j}, 1, \frac{\zeta_{j+1}}{\zeta_j}, \dots, \frac{\zeta_n}{\zeta_j} \right),$$

we have the local coordinate on U_j

$$\left(\frac{\zeta_0}{\zeta_j}, \dots, \frac{\zeta_{j-1}}{\zeta_j}, \frac{\zeta_{j+1}}{\zeta_j}, \dots, \frac{\zeta_n}{\zeta_j} \right). \quad (48)$$

However the above definition of $\mathbf{CP}^n$ is not tractable, so we use the well-known expression by projections

$$\mathbf{CP}^n \cong G_1(\mathbf{C}^{N+1}) = \{P \in M(N+1; \mathbf{C}) \mid P^2 = P, P^\dagger = P \text{ and } \text{tr} P = 1\} \quad (49)$$

and the correspondence

$$[\zeta_0 : \zeta_1 : \cdots : \zeta_n] \longleftrightarrow \frac{1}{|\zeta_0|^2 + |\zeta_1|^2 + \cdots + |\zeta_n|^2} \begin{pmatrix} |\zeta_0|^2 & \zeta_0 \bar{\zeta}_1 & \cdot & \cdot & \zeta_0 \bar{\zeta}_n \\ \zeta_1 \bar{\zeta}_0 & |\zeta_1|^2 & \cdot & \cdot & \zeta_1 \bar{\zeta}_n \\ \cdot & \cdot & & & \cdot \\ \cdot & \cdot & & & \cdot \\ \zeta_n \bar{\zeta}_0 & \zeta_n \bar{\zeta}_1 & \cdot & \cdot & |\zeta_n|^2 \end{pmatrix} \equiv P. \quad (50)$$

If we set

$$|\zeta\rangle = \frac{1}{\sqrt{\sum_{j=0}^n |\zeta_j|^2}} \begin{pmatrix} \zeta_0 \\ \zeta_1 \\ \cdot \\ \cdot \\ \zeta_n \end{pmatrix}, \quad (51)$$

then we can write the right hand side of (50) as

$$P = |\zeta\rangle\langle\zeta| \quad \text{and} \quad \langle\zeta|\zeta\rangle = 1. \quad (52)$$

For example on U_0

$$(z_1, z_2, \dots, z_n) = \left(\frac{\zeta_1}{\zeta_0}, \frac{\zeta_2}{\zeta_0}, \dots, \frac{\zeta_n}{\zeta_0} \right),$$

we have

$$\begin{aligned}
P(z_1, \dots, z_n) &= \frac{1}{1 + \sum_{j=1}^n |z_j|^2} \begin{pmatrix} 1 & \bar{z}_1 & \cdot & \cdot & \bar{z}_n \\ z_1 & |z_1|^2 & \cdot & \cdot & z_1 \bar{z}_n \\ \cdot & \cdot & & & \cdot \\ \cdot & \cdot & & & \cdot \\ z_n & z_n \bar{z}_1 & \cdot & \cdot & |z_n|^2 \end{pmatrix} \\
&= |(z_1, z_2, \dots, z_n)\rangle \langle (z_1, z_2, \dots, z_n)|, \tag{53}
\end{aligned}$$

where

$$|(z_1, z_2, \dots, z_n)\rangle = \frac{1}{\sqrt{1 + \sum_{j=1}^n |z_j|^2}} \begin{pmatrix} 1 \\ z_1 \\ \cdot \\ \cdot \\ z_n \end{pmatrix}. \tag{54}$$

To be clearer, let us give a detailed description for the case of $n = 1$ and 2.

(a) $n = 1$:

$$\begin{aligned}
P(z) &= \frac{1}{1 + |z|^2} \begin{pmatrix} 1 & \bar{z} \\ z & |z|^2 \end{pmatrix} = |z\rangle \langle z|, \\
\text{where } |z\rangle &= \frac{1}{\sqrt{1 + |z|^2}} \begin{pmatrix} 1 \\ z \end{pmatrix}, \quad z = \frac{\zeta_1}{\zeta_0}, \quad \text{on } U_0, \tag{55}
\end{aligned}$$

$$\begin{aligned}
P(w) &= \frac{1}{|w|^2 + 1} \begin{pmatrix} |w|^2 & w \\ \bar{w} & 1 \end{pmatrix} = |w\rangle \langle w|, \\
\text{where } |w\rangle &= \frac{1}{\sqrt{|w|^2 + 1}} \begin{pmatrix} w \\ 1 \end{pmatrix}, \quad w = \frac{\zeta_0}{\zeta_1}, \quad \text{on } U_1. \tag{56}
\end{aligned}$$

(b) $n = 2$:

$$P(z_1, z_2) = \frac{1}{1 + |z_1|^2 + |z_2|^2} \begin{pmatrix} 1 & \bar{z}_1 & \bar{z}_2 \\ z_1 & |z_1|^2 & z_1 \bar{z}_2 \\ z_2 & z_2 \bar{z}_1 & |z_2|^2 \end{pmatrix} = |(z_1, z_2)\rangle \langle (z_1, z_2)|,$$

$$\text{where } |(z_1, z_2)\rangle = \frac{1}{\sqrt{1 + |z_1|^2 + |z_2|^2}} \begin{pmatrix} 1 \\ z_1 \\ z_2 \end{pmatrix}, \quad (z_1, z_2) = \left(\frac{\zeta_1}{\zeta_0}, \frac{\zeta_2}{\zeta_0} \right) \quad \text{on } U_0, \quad (57)$$

$$P(w_1, w_2) = \frac{1}{|w_1|^2 + 1 + |w_2|^2} \begin{pmatrix} |w_1|^2 & w_1 & w_1 \bar{w}_2 \\ \bar{w}_1 & 1 & \bar{w}_2 \\ w_2 \bar{w}_1 & w_2 & |w_2|^2 \end{pmatrix} = |(w_1, w_2)\rangle \langle (w_1, w_2)|,$$

$$\text{where } |(w_1, w_2)\rangle = \frac{1}{\sqrt{|w_1|^2 + 1 + |w_2|^2}} \begin{pmatrix} w_1 \\ 1 \\ w_2 \end{pmatrix}, \quad (w_1, w_2) = \left(\frac{\zeta_0}{\zeta_1}, \frac{\zeta_2}{\zeta_1} \right) \quad \text{on } U_1, \quad (58)$$

$$P(v_1, v_2) = \frac{1}{|v_1|^2 + |v_2|^2 + 1} \begin{pmatrix} |v_1|^2 & v_1 \bar{v}_2 & v_1 \\ v_2 \bar{v}_1 & |v_2|^2 & v_2 \\ \bar{v}_1 & \bar{v}_2 & 1 \end{pmatrix} = |(v_1, v_2)\rangle \langle (v_1, v_2)|,$$

$$\text{where } |(v_1, v_2)\rangle = \frac{1}{\sqrt{|v_1|^2 + |v_2|^2 + 1}} \begin{pmatrix} v_1 \\ v_2 \\ 1 \end{pmatrix}, \quad (v_1, v_2) = \left(\frac{\zeta_0}{\zeta_2}, \frac{\zeta_1}{\zeta_2} \right) \quad \text{on } U_2. \quad (59)$$

B Local Coordinate of the Projector

We give a proof to the last formula in (40).

By making use of the expression by Oike in [14] (we don't repeat it here)

$$\mathcal{P}(\mathcal{Z}) = \begin{pmatrix} \mathbf{1} & -\mathcal{Z}^\dagger \\ \mathcal{Z} & \mathbf{1} \end{pmatrix} \begin{pmatrix} \mathbf{1} & \\ & \mathbf{0} \end{pmatrix} \begin{pmatrix} \mathbf{1} & -\mathcal{Z}^\dagger \\ \mathcal{Z} & \mathbf{1} \end{pmatrix}^{-1} \quad (60)$$

where $\mathcal{Z}$ is some operator on the Fock space $\mathcal{F}$. Let us rewrite this into more useful form. From the simple relation

$$\begin{pmatrix} \mathbf{1} & \mathcal{Z}^\dagger \\ -\mathcal{Z} & \mathbf{1} \end{pmatrix} \begin{pmatrix} \mathbf{1} & -\mathcal{Z}^\dagger \\ \mathcal{Z} & \mathbf{1} \end{pmatrix} = \begin{pmatrix} \mathbf{1} + \mathcal{Z}^\dagger \mathcal{Z} & \\ & \mathbf{1} + \mathcal{Z} \mathcal{Z}^\dagger \end{pmatrix}$$

we have

$$\begin{pmatrix} \mathbf{1} & -\mathcal{Z}^\dagger \\ \mathcal{Z} & \mathbf{1} \end{pmatrix}^{-1} = \begin{pmatrix} (\mathbf{1} + \mathcal{Z}^\dagger \mathcal{Z})^{-1} & \\ & (\mathbf{1} + \mathcal{Z} \mathcal{Z}^\dagger)^{-1} \end{pmatrix} \begin{pmatrix} \mathbf{1} & \mathcal{Z}^\dagger \\ -\mathcal{Z} & \mathbf{1} \end{pmatrix}.$$

Inserting this into (60) and some calculation leads to

$$\mathcal{P}(\mathcal{Z}) = \begin{pmatrix} (\mathbf{1} + \mathcal{Z}^\dagger \mathcal{Z})^{-1} & (\mathbf{1} + \mathcal{Z}^\dagger \mathcal{Z})^{-1} \mathcal{Z}^\dagger \\ \mathcal{Z}(\mathbf{1} + \mathcal{Z}^\dagger \mathcal{Z})^{-1} & \mathcal{Z}(\mathbf{1} + \mathcal{Z}^\dagger \mathcal{Z})^{-1} \mathcal{Z}^\dagger \end{pmatrix}. \quad (61)$$

Comparing (61) with (27) we obtain the “local coordinate”

$$\mathcal{Z} = \frac{1}{R(N) + \theta} a^\dagger = a^\dagger \frac{1}{R(N+1) + \theta} \quad (62)$$

where $R(N) = \sqrt{N + \theta^2}$. $\mathcal{Z}$ obtained by “stereographic projection” is a kind of complex coordinate.

Now if we take a classical limit $a \longrightarrow x - iy$, $a^\dagger \longrightarrow x + iy$ and $\theta = z$ then

$$Z_c = \frac{x + iy}{r + z} \quad (63)$$

where $r = \sqrt{x^2 + y^2 + z^2}$. This is nothing but a well-known one for (10).

C Difficulty of Tensor Decomposition

We point out a difficulty in obtaining the formula (45) or (46) by decomposing tensor products of V .

To obtain the formula (43) there is another method which uses a decomposition of the tensor product $A \otimes A$. Let us introduce. For

$$A = \begin{pmatrix} \alpha & -\bar{\beta} \\ \beta & \bar{\alpha} \end{pmatrix} \in SU(2)$$

we have

$$A \otimes A = \begin{pmatrix} \alpha^2 & -\alpha\bar{\beta} & -\alpha\bar{\beta} & \bar{\beta}^2 \\ \alpha\beta & |\alpha|^2 & -|\beta|^2 & -\bar{\alpha}\bar{\beta} \\ \alpha\beta & -|\beta|^2 & |\alpha|^2 & -\bar{\alpha}\bar{\beta} \\ \beta^2 & \bar{\alpha}\beta & \bar{\alpha}\beta & \bar{\alpha}^2 \end{pmatrix}.$$

For the matrix T coming from the Clebsch–Gordan decomposition

$$T = \begin{pmatrix} 0 & 1 & 0 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} & 0 \\ -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

it is easy to see

$$T^\dagger(A \otimes A)T = \begin{pmatrix} |\alpha|^2 + |\beta|^2 & & & \\ & \alpha^2 & -\sqrt{2}\alpha\bar{\beta} & \bar{\beta}^2 \\ & \sqrt{2}\alpha\beta & |\alpha|^2 - |\beta|^2 & -\sqrt{2}\bar{\alpha}\bar{\beta} \\ & \beta^2 & \sqrt{2}\bar{\alpha}\beta & \bar{\alpha}^2 \end{pmatrix} = \begin{pmatrix} 1 & \\ & \phi_1(A) \end{pmatrix} \quad (64)$$

where we have used $|\alpha|^2 + |\beta|^2 = 1$. This means a well-known decomposition

$$\frac{1}{2} \otimes \frac{1}{2} = 0 \oplus 1.$$

Let us take an analogy. For

$$V = \begin{pmatrix} X_0 & -Y_0^\dagger \\ Y_0 & X_{-1} \end{pmatrix}$$

we have

$$V \otimes V = \begin{pmatrix} X_0^2 & -X_0Y_0^\dagger & -Y_0^\dagger X_0 & Y_0^\dagger Y_0^\dagger \\ X_0Y_0 & X_0X_{-1} & -Y_0^\dagger Y_0 & -Y_0^\dagger X_{-1} \\ Y_0X_0 & -Y_0Y_0^\dagger & X_{-1}X_0 & -X_{-1}Y_0^\dagger \\ Y_0Y_0 & Y_0X_{-1} & X_{-1}Y_0 & X_{-1}^2 \end{pmatrix}.$$

However, the analogy breaks down at this stage because of the non-commutativity

$$T^\dagger(V \otimes V)T \neq \begin{pmatrix} \mathbf{1} & \\ & \Phi_1(V) \end{pmatrix} \quad (65)$$

for (45). We leave it to the readers. There is no (well-known) direct method to obtain $\Phi_1(V)$ at the current time.

Last, let us make a comment. For the matrix T coming from the Clebsch–Gordan decomposition (see [10])

$$T = \begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{6}} & 0 & 0 & \frac{1}{\sqrt{3}} & 0 & 0 \\ -\frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{6}} & 0 & 0 & \frac{1}{\sqrt{3}} & 0 & 0 \\ 0 & 0 & 0 & \frac{\sqrt{2}}{\sqrt{3}} & 0 & 0 & \frac{1}{\sqrt{3}} & 0 \\ 0 & 0 & -\frac{\sqrt{2}}{\sqrt{3}} & 0 & 0 & \frac{1}{\sqrt{3}} & 0 & 0 \\ 0 & \frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{6}} & 0 & 0 & \frac{1}{\sqrt{3}} & 0 \\ 0 & -\frac{1}{\sqrt{2}} & 0 & -\frac{1}{\sqrt{6}} & 0 & 0 & \frac{1}{\sqrt{3}} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

it is not difficult to see

$$\begin{aligned} & T^\dagger(A \otimes A \otimes A)T \\ &= \begin{pmatrix} \alpha & -\bar{\beta} & & & & & & \\ \beta & \bar{\alpha} & & & & & & \\ & & \alpha & -\bar{\beta} & & & & \\ & & \beta & \bar{\alpha} & & & & \\ & & & & \alpha^3 & -\sqrt{3}\alpha^2\bar{\beta} & \sqrt{3}\alpha\bar{\beta}^2 & -\bar{\beta}^3 \\ & & & & \sqrt{3}\alpha^2\beta & (|\alpha|^2 - 2|\beta|^2)\alpha & -(2|\alpha|^2 - |\beta|^2)\bar{\beta} & \sqrt{3}\bar{\alpha}\bar{\beta}^2 \\ & & & & \sqrt{3}\alpha\beta^2 & (2|\alpha|^2 - |\beta|^2)\beta & (|\alpha|^2 - 2|\beta|^2)\bar{\alpha} & -\sqrt{3}\bar{\alpha}^2\bar{\beta} \\ & & & & \beta^3 & \sqrt{3}\bar{\alpha}\beta^2 & \sqrt{3}\bar{\alpha}^2\beta & \bar{\alpha}^3 \end{pmatrix} \\ &= \begin{pmatrix} \phi_{1/2}(A) & & & \\ & \phi_{1/2}(A) & & \\ & & \phi_{3/2}(A) & \end{pmatrix}. \end{aligned} \tag{66}$$

This means a well-known decomposition

$$\frac{1}{2} \otimes \frac{1}{2} \otimes \frac{1}{2} = (0 \oplus 1) \otimes \frac{1}{2} = \left(0 \otimes \frac{1}{2}\right) \oplus \left(1 \otimes \frac{1}{2}\right) = \frac{1}{2} \oplus \frac{1}{2} \oplus \frac{3}{2}.$$

References

- [1] K. Fujii : Jaynes–Cummings Model and a Non–Commutative “Geometry” : A Few Problems Noted, quant-ph/0410201.
- [2] M. Nakahara : Geometry, Topology and Physics, Adam Hilger, Bristol and New York, 1990.
- [3] R. Rajaraman : Solitons and Instantons, North–Holland, Amsterdam, 1982.
- [4] K. Fujii : Classical solutions of higher dimensional non–linear sigma models on spheres, Lett. Math. Phys. 10(1985), 49 ; K. Fujii : Extended Yang–Mills Models on Even Dimensional Spaces, Lett. Math. Phys. 12(1986), 363 ; B. Grossman, T. W. Kephart and J. D. Stasheff : Solutions to Yang–Mills Field Equations in Eight–Dimensions and the Last Hopf Map, Commun. Math. Phys. 96(1984), 431, and 100(1985), 311 ; K. Fujii, S. Kitakado and Y. Ohnuki : Gauge Structure on S^D , Mod. Phys. Lett. 10(1995), 867.
- [5] A. Shapere and F. Wilczek : Geometric Phases in Physics, World Scientific, 1989.
- [6] A. P. Balachandran and G. Immirzi : Fuzzy Nambu–Goldstone Physics, Int. J. Mod. Phys. A 18(2003), 5931, hep-th/0212133.
- [7] G. Sardanashvily and G. Giachetta : What is Geometry in Quantum Theory, Int. J. Geom. Methods Mod. Phys. 1(2004), 1, hep-th/0401080.
- [8] E. T. Jaynes and F. W. Cummings : Comparison of Quantum and Semiclassical Radiation Theories with Applications to the Beam Maser, Proc. IEEE 51(1963), 89 ; M. Tavis and F. W. Cummings : Exact Solution for an N–Molecule–Radiation–Field Hamiltonian, Phys. Rev. 170(1968), 379 ; R. H. Dicke : Coherence in Spontaneous Radiation Processes, Phys. Rev, 93(1954), 99.
- [9] P. Meystre and M. Sargent III : Elements of Quantum Optics (third edition), Springer–Verlag, 1990 ;

- [10] K. Fujii, K. Higashida, R. Kato and Y. Wada : Explicit Form of the Solution of Two Atoms Tavis–Cummings Model, to appear in The Bulletin of Yokohama City University, vol.56(2005), quant-ph/0403008 ; K. Fujii, K. Higashida, R. Kato, T. Suzuki and Y. Wada : Explicit Form of the Evolution Operator of Tavis-Cummings Model : Three and Four Atoms Cases, Int. J. Geom. Methods Mod. Phys, vol.1(2004), 721, quant-ph/0409068.
- [11] K. Fujii, K. Higashida, R. Kato and Y. Wada : Cavity QED and Quantum Computation in the Weak Coupling Regime, J. Opt. B: Quantum and Semiclass. Opt, 6(2004) 502, quant-ph/0407014 ; K. Fujii, K. Higashida, R. Kato and Y. Wada : Cavity QED and Quantum Computation in the Weak Coupling Regime II : Complete Construction of the Controlled–Controlled NOT Gate, Contribution to the volume “Quantum Computing: New Research” to be published by Nova Science Publishers (USA), quant-ph/0501046.
- [12] K. Fujii, K. Higashida, R. Kato, T. Suzuki and Y. Wada : Quantum Diagonalization Method in the Tavis–Cummings Model, to appear in Int. J. Geom. Methods Mod. Phys, vol.2, no.3(2005), quant-ph/0410003.
- [13] T. Suzuki : in preparation.
- [14] K. Fujii : Introduction to Grassmann Manifolds and Quantum Computation, J. Applied Math, 2(2002), 371, quant-ph/0103011.
- [15] K. Funahashi, T. Kashiwa, S. Sakoda and K. Fujii : Coherent states, path integral, and semiclassical approximation, J. Math. Phys., 36(1995), 3232 ; K. Funahashi, T. Kashiwa, S. Sakoda and K. Fujii : Exactness in the Wentzel-Kramers-Brillouin approximation for some homogeneous spaces, J. Math. Phys., 36(1995), 4590 ; K. Fujii, T. Kashiwa, S. Sakoda : Coherent states over Grassmann manifolds and the WKB exactness in path integral, J. Math. Phys., 37(1996), 567.
- [16] A. Asada : Regularized Calculus : An Application of Zeta Regularization to Infinite Dimensional Geometry and Analysis, Int. J. Geom. Methods Mod. Phys, vol.1(2004), 107.